

Math 254-2 Exam 5 Solutions

1. Carefully state the definition of “basis”. Give two examples from \mathbb{R}^2 .

A basis is a set of vectors that is both independent and spanning. Equivalently, a basis is a maximal set of independent vectors. Equivalently, a basis is a minimal set of spanning vectors. Many examples are possible, such as $\{(1, 0), (0, 1)\}$ or $\{(1, 1), (2, 3)\}$. All must contain exactly two, linearly independent, vectors.

Problems 2 and 3 both concern the matrix $A = \begin{pmatrix} 2 & -4 & 6 & 0 & 4 \\ 1 & -2 & 3 & 0 & 2 \\ -1 & 2 & -3 & 1 & -1 \\ -2 & 4 & -6 & 2 & -2 \\ 3 & -6 & 9 & -3 & 3 \end{pmatrix}$.

2. Set $S = \text{Rowspace}(A)$. Find a basis for S , and determine its dimension.

$\begin{pmatrix} 2 & -4 & 6 & 0 & 4 \\ 1 & -2 & 3 & 0 & 2 \\ -1 & 2 & -3 & 1 & -1 \\ -2 & 4 & -6 & 2 & -2 \\ 3 & -6 & 9 & -3 & 3 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 3 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & -3 & -3 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & -2 & 3 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$, in row echelon form. There are two pivots, hence S is two dimensional. Many bases are possible; the natural one is the nonzero rows of the echelon matrix: $\{(1, -2, 3, 0, 2), (0, 0, 0, 1, 1)\}$. However any two independent elements of the rowspace would also work, such as two independent rows of A itself: $\{(2, -4, 6, 0, 4), (-1, 2, -3, 1, -1)\}$. The first two rows of A will NOT work, since they are dependent.

3. Set $T = \text{Columnspace}(A)$. Find a basis for T , and determine its dimension.

The rowspace and columnspace have the same dimension, hence T is two dimensional. The pivots of the row echelon form of A are in the first and fourth columns, hence the first and fourth columns of A form a basis for the columnspace: $\{(2, 1, -1, -2, 3)^T, (0, 0, 1, 2, -3)^T\}$. This is not the only basis; any two independent elements of the columnspace would also work.

Problems 4 and 5 both concern the vector spaces $A = \text{Span}((2, 0, 1), (1, -1, 3))$ and $B = \text{Span}((5, 1, 0), (0, 4, -10))$. Both are subspaces of \mathbb{R}^3 .

4. Find a basis for $A + B$, and determine its dimension.

We begin by putting the generating vectors in a matrix, then putting this matrix into echelon form: $\begin{pmatrix} 2 & 1 & 5 & 0 \\ 0 & -1 & 1 & 4 \\ 1 & 3 & 0 & -10 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & -5 & 5 & 20 \\ 0 & -1 & 1 & 4 \\ 1 & 3 & 0 & -10 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 3 & 0 & -10 \\ 0 & -1 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{pmatrix}$. This has two pivots, hence $A + B$ is two dimensional. The pivots are in the first two columns, hence a basis is $\{(2, 0, 1), (1, -1, 3)\}$ (other bases are possible). Note: this solution has the matrix as 3×4 , putting the vectors into columns. It is equally correct to put the vectors into rows, giving a 4×3 matrix.

5. Find a basis for $A \cap B$, and determine its dimension.

$\dim(A + B) + \dim(A \cap B) = \dim(A) + \dim(B)$. We have $\dim(A) = \dim(B) = \dim(A + B) = 2$, hence we can solve and determine $\dim(A \cap B) = 2$. Hence $A \cap B = A = B = A + B$, so a basis for $A \cap B$ is any basis for A (or B), such as $\{(2, 0, 1), (1, -1, 3)\}$.