

## Math 254-2 Exam 9 Solutions

1. Carefully define the term “spanning”. Give two examples in  $\mathbb{R}^2$ .

A set of vectors is spanning if every vector in the vector space can be expressed as a linear combination of vectors from this set. Many examples are possible, e.g.  $\{(1, 0), (0, 1)\}$ ,  $\{(1, 0), (0, 1), (1, 1), (2, 3)\}$ .

2. Consider the basis  $S = \{(1, 2), (2, 5)\}$  of  $\mathbb{R}^2$ , and the linear operator  $F(x, y) = (2x - 3y, x - y)$ . Find the matrix representation  $[F]_S$ .

We have  $P_{ES} = \begin{pmatrix} 1 & 2 \\ 2 & 5 \end{pmatrix}$ , so  $P_{SE} = P_{ES}^{-1} = \begin{pmatrix} 5 & -2 \\ -2 & 1 \end{pmatrix}$ . We calculate  $[F]_E = ([F(e_1)]_E \ [F(e_2)]_E) = \left( \begin{bmatrix} 2 \\ 1 \end{bmatrix}_E \ \begin{bmatrix} -3 \\ -1 \end{bmatrix}_E \right) = \begin{pmatrix} 2 & -3 \\ 1 & -1 \end{pmatrix}$ . Hence  $[F]_S = P_{SE}[F]_E P_{ES} = \begin{pmatrix} 5 & -2 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} 2 & -3 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 2 & 5 \end{pmatrix} = \begin{pmatrix} -18 & -49 \\ 7 & 19 \end{pmatrix}$ .

3. Prove that, for any square matrices  $A, B$ , if  $A$  is similar to  $B$ , then  $B$  must be similar to  $A$ .

Suppose that  $A$  is similar to  $B$ . Then there is some invertible matrix  $P$  with  $A = PBP^{-1}$ . Multiply this expression on the left by  $P^{-1}$ , and on the right by  $P$ , to get  $P^{-1}AP = P^{-1}PBP^{-1}P = IBI = B$ . Hence, there is some invertible matrix  $Q = P^{-1}$ , such that  $B = QAQ^{-1}$ , so  $B$  is similar to  $A$ .

For the last two questions, set  $V$  to be the vector space of functions that have as a basis  $S = \{1, \sin \theta, \cos \theta, \sin 5\theta, \cos 5\theta\}$ .

4. Let  $D$  be the differential operator on  $V$ ,  $D(f(\theta)) = f'(\theta)$ . Find the matrix representation  $[D]_S$ .

$$[D]_S = ([D(1)]_S \ [D(\sin \theta)]_S \ [D(\cos \theta)]_S \ [D(\sin 5\theta)]_S \ [D(\cos 5\theta)]_S) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -5 \\ 0 & 0 & 0 & 5 & 0 \end{pmatrix}.$$

5. Let  $L$  be the operator on  $V$  given by  $L(f(\theta)) = f''(\theta) - 2f(\theta)$ . Find the matrix representation  $[L]_S$ .

$$[L]_S = ([L(1)]_S \ [L(\sin \theta)]_S \ [L(\cos \theta)]_S \ [L(\sin 5\theta)]_S \ [L(\cos 5\theta)]_S) = \begin{pmatrix} -2 & 0 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 & 0 \\ 0 & 0 & -3 & 0 & 0 \\ 0 & 0 & 0 & -27 & 0 \\ 0 & 0 & 0 & 0 & -27 \end{pmatrix}.$$